

SHORT COMMUNICATIONS

J. Raptor Res. 38(4):357–361

© 2004 The Raptor Research Foundation, Inc.

GENDER DETERMINATION IN THE SWAINSON'S HAWK (*BUTEO SWAINSONI*) USING MOLECULAR PROCEDURES AND DISCRIMINANT FUNCTION ANALYSIS

JOSÉ HERNÁN SARASOLA¹ AND JUAN JOSÉ NEGRO

Department of Applied Biology, Estación Biológica de Doñana (CSIC), Avda. de María Luisa s/n, Pabellón del Perú, 41013 Sevilla, Spain

KEY WORDS: *Swainson's Hawk*; *Buteo swainsoni*; *gender determination*; *molecular sexing*; *morphometric measures*; *wintering grounds*.

Gender identification of individuals is important in many studies of wild animals. However, easy determination of sex is difficult for monomorphic species including most birds of prey. Many raptor species show little plumage dimorphism, and although females are generally larger than males, overlap in morphometric measurements and body mass make gender determination difficult even when birds are captured and handled.

The Swainson's Hawk (*Buteo swainsoni*) breeds in North America and migrates to southern South America during the boreal winter that involves a trip of ca. 10 000 km each way (second longest migration distance among raptors; England et al. 1997, Fuller et al. 1998). Male and female Swainson's Hawks are similar in plumage when adults (Wheeler and Clark 1995, England et al. 1997), and as in most of the genus *Buteo*, plumage polymorphism occurs both in immature and adult birds. Although accurate methods for gender determination using morphometric data have been developed for several raptor species (Bortolotti 1984a, 1984b, Garcelon et al. 1985, Edwards and Kochert 1986, Ferrer and De Le Court 1992, Balbontín et al. 2001, Palma et al. 2001), no reliable criteria for gender determination of Swainson's Hawks using external characteristics have been described.

During the last decade, the development of laboratory techniques involving molecular procedures has provided reliable methods for the accurate gender determination of the majority of avian species (Ellegren and Sheldon 1997). PCR-based methods targeting CHD1-Z and CHD1-W genes are purported to be of universal application to birds, with the exception of ratite species (Ellegren 1996, Fridolfsson and Ellegren 1999). The aim of this study was to develop an accurate method for gender determination of Swainson's Hawks using molecular procedures and

morphometric criteria. Our goal was to obtain a general model, derived from discriminant analysis, to determine gender of immature and adult Swainson's Hawks.

METHODS

We captured and sampled free-living Swainson's Hawks during two wintering seasons (austral summers) in three study areas located in central Argentina. We captured 34 hawks in the vicinity of a roost site near Las Varillas, Córdoba province (31°58'S, 62°50'W), from 19–26 January 2003. One hawk was captured in northern La Pampa province (35°14'S, 63°57'W) on 21 November 2002, and 34 hawks from 7–10 December 2003 at the same site. The sample was completed with 35 hawks captured near Santa Rosa (36°33'S, 64°07'W), La Pampa province, from 21–29 January 2004. The habitat where trapping was conducted consisted of agricultural fields of continuous crops, with soybeans as the principal crop. Planted pastures and natural fields comprised the remaining habitat.

Hawks were captured in open fields near the roost using bal-chatri traps (Berger and Mueller 1959) in early morning and during the afternoon. Traps were set in front of fence posts usually used by hawks for perching, both when they left roosts in the morning and during late afternoon before returning. Captured hawks were aged as juveniles or adults based on plumage characteristics (Wheeler and Clark 1995), with immature birds grouped with juveniles using the same criteria employed by Goldstein et al. (1999). Hawks were banded and weighed with a 1500 g Pesola scale (Pesola AG, Baar, Switzerland) to the nearest 2 g. Six morphometric measurements were taken from adults and juveniles. We measured the length of wing chord (WING) and tail (TAIL) using a plastic rule to the nearest 1 mm, and length of the exposed culmen (CULMEN), tarsus (TARSUS), and hallux claw (HALLUX) using a caliper to the nearest 0.05 mm. We also measured the forearm length (FORE-ARM), or the length from the front of the folded wrist to the proximal extremity of the ulna, also using a caliper (Ferrer and De Le Court 1992, Balbontín et al. 2001; Fig 1). For a few birds only some of the body measurements were recorded (Tables 1, 2).

Approximately 2 ml of blood were taken from each bird from the brachial vein. The blood was placed in tubes with 96% ethanol that were kept in coolers until analysis in the laboratory. The cellular fraction of the

¹ E-mail address: sarasola@ebd.csic.es

Table 2. Morphometric measurements of Swainson’s Hawks in wintering grounds and analysis of variance (ANOVA) test results for gender differences. All measurements except mass (g) are in mm.

	FEMALES				MALES				ANOVA	
	\bar{x}	SD	RANGE	N	\bar{x}	SD	RANGE	N	F	P
Wing	409.0	11.0	390.0–430.0	38	383.0	11.0	350.0–420.0	66	127.74	<0.001
Tail	211.0	9.0	180.0–240.0	38	195.0	8.0	180.0–230.0	66	88.98	<0.001
Culmen	24.1	1.1	21.0–26.1	37	22.3	1.0	20.3–25.6	65	69.53	<0.001
Tarsus	72.6	4.0	64.6–80.1	37	69.5	3.8	60.3–78.4	66	15.68	<0.001
Hallux	25.3	1.1	22.6–28.1	38	23.7	1.0	21.4–26.4	66	55.97	<0.001
Forearm	144.2	4.8	134.5–157.0	38	134.0	4.3	124.0–144.8	66	122.68	<0.001
Mass	895.1	118.3	590.0–1110.0	38	735.6	66.8	540.0–880.0	65	76.64	<0.001

$D < 0$ identified males. The dividing point between genders for forearm length obtained by solving for 0 was 140.2 mm, with values over this point representing females and values under it representing males. Forearm, tail length, and wing chord length were retained in the stepwise discriminant analysis. The resulting linear function ($D_2 = 0.36 \text{ FOREARM} + 1.36 \text{ TAIL} + 1.04 \text{ WING} - 120.95$) increased our predictive power against D_1 (Table 3) and a lower value of Wilk’s lambda indicated that females and males were better separated with this linear combination of variables than using only forearm.

DISCUSSION

Our data indicated that there were significant differences between male and female Swainson’s Hawks. However, there was considerable overlap in the ranges of the morphometric measurements, suggesting that the use of relative size as the only criterion to determine gender in this raptor would result in errors.

Bill depth, toe-pad length, and body mass are morphometric measurements frequently used in determining gender of size-dimorphic birds of prey (Bortollotti 1984a,

1984b, Garcelon et al. 1985, Edwards and Kochert 1986). Forearm length is a body feature traditionally not recorded for wild raptors, but its use has become more common as it is a measure easily obtained from museum skins. Furthermore, it has been shown to be a low variance measure with high repeatability among observers (Ferrer and De Le Court 1992). These features make forearm a reliable morphometric measure that should be considered as a general and standard measure in morphometric gender determination of birds of prey.

Ideally, a technique for gender determination would be applicable to individuals of all ages and under different conditions. Our model based on forearm as a single explanatory variable can be used even on birds showing incomplete molt or evidence of loss of corporal mass due to fasting. The alternative model (D_2) can be also applied in cases in which loss or gain of body mass is suspected to occur (see Smith et al. 1986, Goldstein et al. 1999). The second function, which classified the highest percentage of cases after the cross-validation and chance correction tests, produced a better separation of groups (lower value of Wilk’s lambda), but needed more vari-




Figure 2. Gender determination using PCR methods. A multiple amplification with 2550F amplify a 420 bp fragment of W chromosome in females and 2550F + 2718R that amplifies 600 bp fragments in both sexes. Males and females are indicated as M and F, respectively.

Table 3. Accuracy of sexing Swainson's Hawks obtained from discriminant analysis using single measurements or combinations of morphometric variables, and assessed by cross-validation procedure and Cohen's Kappa calculation. Single variables are ordered from higher to lower values of Cohen's Kappa.

VARIABLE	CASES CORRECTLY SEPARATED					COHEN'S KAPPA	P
	WILK'S LAMBDA	N	PERCENT FEMALES	PERCENT MALES	PERCENT OVERALL		
Forearm	0.454	(104)	81.6	93.9	89.4	0.77	<0.01
Wing	0.444	(104)	81.6	90.9	87.5	0.73	<0.01
Tail	0.534	(104)	81.6	90.9	87.5	0.73	<0.01
Tarsus	0.866	(103)	70.0	87.5	82.4	0.28	>0.05
Forearm + Tail + Wing	0.332	(104)	86.8	97.0	93.3	0.85	<0.01

ables to be applicable. Its use would be better when complete data sets of the variables are available and given that measurements are not biased by external factors (e.g., molting). Furthermore, D_1 discriminant function allows for gender determination of dead birds when natural decomposition and the effect of scavengers after days of exposure in the field makes classification of sex by standard forensic methods impossible.

Insecticide poisoning of hawks in their wintering ground during 1995–96 and 1996–97 austral summers (Woodbridge et al. 1995, Goldstein et al. 1996) has been documented in this species, with ca. 20 000 birds poisoned in 1996–97. Gender determination of Swainson's Hawks would provide a valuable tool for a complete assessment of these mortality incidents, including the gender of affected birds.

RESUMEN.—*Buteo swainsoni* es un ave de presa poco dimórfica, y aunque las hembras suelen ser más grandes que los machos, la determinación del sexo en esta especie puede ser difícil, aún cuando las aves son capturadas y manipuladas. En este artículo presentamos un método de sexado para *B. swainsoni* basado en técnicas moleculares y análisis discriminantes. Los datos empleados corresponden a medidas morfométricas de 104 individuos silvestres capturados en el área de invernada de la especie durante los veranos australes 2002–03 y 2003–04. Encontramos diferencias significativas entre machos y hembras en todas las medidas morfométricas consideradas, mientras que los juveniles se diferenciaron de los adultos sólo en su masa corporal, la longitud del culmen y la longitud del hálux. Usando sólo la longitud del antebrazo como variable de predicción, nuestra función discriminante clasificó correctamente el 89.4% de los machos y el 93.9% de las hembras. Una segunda función que incluía la longitud del antebrazo, de la cola y de la cuerda alar mejoró la separación de los grupos y también el porcentaje de individuos correctamente clasificados (97.0% y 93.3% de los machos y hembras, respectivamente). El uso de medidas relacionadas con el tamaño estructural de las aves como la longitud del antebrazo,

de la cola y del ala hacen de éste un método seguro y de amplia aplicación, aún para aves pertenecientes a distintas clases de edad.

[Traducción de los autores]

ACKNOWLEDGMENTS

We wish to thank N. Witheman, R.A. Sosa, V. Salvador, and M. Santillán for help in trapping hawks during fieldwork. P. Bazaga did the lab work and J. Balbontín helped with data analysis. We are grateful for comments by J. Balbontín, M. Bechard, and an anonymous reviewer that greatly improved this manuscript. We also thank personnel from Ea. La Independencia and Ea. La Armonía II, the Agencia Córdoba Ambiente of Córdoba province, and the Dirección de Recursos Naturales of La Pampa province for providing permission to carry out trapping in the study areas. We thank A. Lanusse, S. Salva, M. Aliaga, and J. Montoya for their hospitality and logistic support during fieldwork in La Pampa province. J.H. Sarasola had a fellowship from the Consejo Nacional de Investigaciones Científicas y Técnicas of Argentina. This study was supported by a grant from the Wildlife Conservation Society (U.S.A.).

LITERATURE CITED

BALBONTÍN, J., M. FERRER, AND E. CASADO. 2001. Sex determination in Booted Eagles (*Hieraaetus pennatus*) using molecular procedures and discriminant function analysis. *J. Raptor Res.* 35:20–23.

BERGER, D.D. AND H.C. MUELLER. 1959. The bal-chatri: a trap for the birds of prey. *Bird-Banding* 30:18–26.

BORTOLOTTI, G.R. 1984a. Age and sex size variation in Golden Eagles. *J. Field Ornithol.* 55:54–56.

———. 1984b. Criteria for determining age and sex of nestling Bald Eagles. *J. Field Ornithol.* 55:467–481.

EDWARDS, T.C. AND M.N. KOCHIERT. 1986. Use of body weight and length of footpad as predictors of sex in Golden Eagles. *J. Field Ornithol.* 57:317–319.

ELLEGREN, H. 1996. First gene on the avian W chromosome (CHD) provides a tag for universal sexing of non-ratite birds. *Proc. R. Soc. Lond. B.* 263:1635–1641.

——— AND B.C. SHELDON. 1997. New tools for sex iden-

- tification and the study of sex allocation in birds. *Trends Ecol. Evol.* 12:255–259.
- ENGLAND, A.S., M.J. BECHARD, AND C.S. HOUSTON. 1997. Swainson's Hawk (*Buteo swainsoni*). In A. Poole and F. Gill [Eds.], *The birds of North America*, No. 265. The American Ornithologists' Union, Washington, DC U.S.A.
- FERRER, M. AND C. DE LE COURT. 1992. Sex identification in the Spanish Imperial Eagle. *J. Field Ornithol.* 62:359–364.
- FRIDOLFSSON, A.K. AND H. ELLEGREN. 1999. A simple and universal method for molecular sexing of non-ratite birds. *J. Avian Biol.* 30:116–121.
- FULLER, M.R., W.S. SEEGAR, AND L.S. SCHUECK. 1998. Routes and travel rates of migrating Peregrine Falcons *Falco peregrinus* and Swainson's Hawk *Buteo swainsoni* in the Western Hemisphere. *J. Avian Biol.* 29:433–440.
- GARCELON, D.K., M.S. MARTELL, P.T. REDIG, AND L.C. BUEN. 1985. Morphometric, karyotypic, and laparoscopic techniques for determining sex in Bald Eagles. *J. Wildl. Manage.* 49:595–599.
- GOLDSTEIN, M.I., B. WOODBRIDGE, M.E. ZACCAGNINI, S.B. CANAVELLI, AND A. LANUSSE. 1996. An assessment of mortality of Swainson's Hawks on wintering grounds in Argentina. *J. Raptor Res.* 30:106–107.
- , P.H. BLOOM, J.H. SARASOLA, AND T.E. LACHER. 1999. Post-migration weight gain of Swainson's Hawks in Argentina. *Wilson Bull.* 111:428–432.
- MANLY, B.F.J. 1986. *Multivariate statistical methods a primer*. Chapman and Hall, London, U.K.
- PALMA, L., S. MIRA, P. CARDIA, P. BEJA, T. GUILLEMAUD, N. FERRAND, M.L. CANCELA, AND L. CANCELA DA FONSECA. 2001. Sexing Bonelli's Eagle nestlings: morphometrics versus molecular techniques. *J. Raptor Res.* 35:187–193.
- SMITH, N.G., D.L. GOLDSTEIN, AND G.A. BARTHOLOMEW. 1986. Is long-distance migration possible using only stored fat? *Auk* 103:607–611.
- TITUS, K. AND J.A. MOSHER. 1984. Chance-corrected classification for use in discriminant analysis: ecological applications. *Am. Midl. Nat.* 111:1–7.
- WHEELER, B.K. AND W.S. CLARK. 1995. *A photographic guide to North American raptors*. Academic Press, San Diego, CA U.S.A.
- WOODBRIDGE, B., K.K. FINLEY, AND S.T. SEAGER. 1995. An investigation of the Swainson's Hawk in Argentina. *J. Raptor Res.* 29:202–204.
- ZAR, J.H. 1996. *Biostatistical analysis*, 3rd Ed. Prentice Hall, Princeton, NJ U.S.A.

Received 12 August 2003; accepted 11 July 2004

J. Raptor Res. 38(4):361–366

© 2004 The Raptor Research Foundation, Inc.

PRODUCTIVITY AND FLEDGLING SEX RATIO IN A CINEREOUS VULTURE (*AEGYPIUS MONACHUS*) POPULATION IN SPAIN

AUXILIADORA VILLEGAS,¹ JUAN MANUEL SÁNCHEZ-GUZMÁN, EMILIO COSTILLO, CASIMIRO CORBACHO, AND RICARDO MORÁN

Grupo de Investigación en Conservación, Área de Zoología, Universidad de Extremadura, Avda. Elvas s/n, 06071 Badajoz, Spain

KEY WORDS: *Cinereous Vulture*; *Aegypius monachus*; *breeding conditions*; *brood size*; *fledgling-sex ratio*; *sexual dimorphism*; *Spain*.

Upon initial review, the mechanism of chromosomal gender determination in birds and mammals seems to be a factor limiting the parents' ability to modify the sex ratio of their progeny (Charnov 1982). However, sex allocation theory (Fisher 1930, Charnov 1982) predicts that the sex ratio can deviate from the expected 1:1, particularly when the costs of rearing the two genders are

different. The optimal sex allocation for individuals can be predicted from three basic non-mutually exclusive hypotheses (reviewed by Frank 1990). (1) Fisher (1930) proposed that parental expenditure in the population should be equal for all sons and daughters, which would result in a population sex ratio biased toward the gender that costs less to produce. (2) Trivers and Willard (1973) hypothesized that if the reproductive return differs between genders depending on parental condition at the time of breeding, natural selection would favor facultative adjustments of offspring sex ratios to obtain the maximum fitness from a breeding attempt. (3) Charnov (1982) generalized Trivers and Willard's hypothesis to cover any socio-environmental variable that might pre-

¹ E-mail address: villegas@unex.es